

Instructions for use - Please read carefully!

Hospisept®

Ready for use

Alcohol based hand and skin disinfectant

DE-Reg-No: 6339833.00.00

Composition

100 g **solution** contains:

Active substance: 55 g Propan-1-ol (Ph. Eur.), 15.84 g Ethanol 96%

Other substances: Dibutyl adipate, Butan-2-on, Purified Water

Range of application

- Control of bacteria (incl. tb), fungi and enveloped viruses
- Hygienic and surgical hand disinfection,
- pre-surgical skin disinfection, nursing of sutures
- Before intramuscular and intravenous injections as well as punctures

Mode of application / dosage

Use externally and undiluted only!

Bactericidal (incl. tb), fungicidal

virucidal* against enveloped viruses incl. HBV/HCV/HIV

and Rota virus

conc. - 30 sec.

Virucidal*

conc. - 5 min.

Adeno virus

conc. - 1 min.

*according to recommendation of Robert Koch Institute Berlin/ Federal Health Gazette 01/2004

Surgical hand disinfection

Pour 5 ml **Hospisept®** in the cupped hands and rub on hands and forearms until dry giving particular attention to the fingertips and to the areas around the nails**. Duration: 1 minute.

Hygienic hand disinfection

(Also appropriate for use according to § 18 of the Federal Law on Epidemics)

At least 3 ml during 30 seconds**.

**Make sure that the skin is wet with the disinfectant during the complete disinfecting time

Skin disinfection

Before punctures and injections:

conc. at least 15 sec

Before punctures of articulations, body cavities, hollow organs

and before surgery: Wet the area to be disinfected completely with **Hospisept®** and allow to dry.

Exposure time:

conc. at least 1 min;

before all medical/surgical interventions on areas rich in sebaceous glands (e.g. on the forehead):

conc. at least 1 min.

Contra-indications:

Hypersensitivity against components.

Interactions

None known!

Side effects

Especially by frequent use the skin might become dry and irritated (reddening, burning feeling). Allergic reactions are very seldom. To prevent extreme skin dryness we recommend to use skin cream during breaks and at the end of work. If skin gets very sore consult a physician.

"Reporting of side-effects:

If you get any side-effects, talk to your doctor/ pharmacist or healthcare professional. This also includes any possible side effects not listed in this leaflet. You can also report side effects directly to the Bundesinstitut für Arzneimittel und Medizinprodukte, Abt. Pharmakovigilanz, Kurt-Georg-Kiesinger Allee 3, D-53175 Bonn, or on their homepage <http://www.bfarm.de>. By reporting side effects you can help provide more information on the safety of this medicinal product."

Warnings

a) For the application of **Hospisept®** on children's skin, during pregnancy and lactation sufficient investigations are not available. In such cases a physician or a pharmacist should be asked for advice before application. In the case of babies and small children apply on large skin surfaces only under medical supervision. No other side effects as mentioned are to be expected when used according to recommendations of manufacturer.

b) Do not apply on **open** wounds!

If **Hospisept®** has come into contact with mucous membranes like **eyes**, throat, stomach, bowels etc., if applied on open wounds, rinse thoroughly with **plenty** of water and get medical attention immediately.

c) If **Hospisept®** has been taken **orally**, make patient **drink plenty** of water to dilute the product and get medical attention immediately.

Safety advice for storage:

Keep medicinal drugs out of reach of children, not above 25°C and away from ignition sources. Inflammable (flash point acc. to DIN 51 755: 26 °C). Do not spray into open flame! Do not use beyond expiry date! Shelf-life (including after opening bottle): see label/bottle.

Pharmaceutical Data

- Transfer product into other containers only under aseptic conditions (for ex. "clean bench").
- Shelf-life: After opening the container: up to the expiry date. If the product is used in a wall dispenser: 6 months but not beyond the expiry date.

Use drugs safely. Always read the label and product information before use.

Properties

- Comprehensive range of action
- Good cleaning power and reliable action even in presence of organic substances like albumen, serum and blood
- Well tolerated by the skin, even when used frequently

We recommend using the following compatible products before proceeding to the surgical disinfection of the hands and after finishing work:

For cleaning: **Majola®**, **Wasa® 2000** or **Wasa-Soft®**

For care/protection: **Majola® - H5 Cream**

Application schedule for hygienic hand disinfection

Hygienic hand disinfection	Product	Way of application
Common situation: Disinfection	Hospisept®	Pour 3 ml Hospisept® in the cupped hands and rub until dry. Duration: ½ min
Dirty hands:		(Take off dirt with tissue soaked with Hospisept® and eliminate dirt - then disinfect as described above.)

Disinfection then Cleaning	Hospisept® Majola® or Wasa® 2000	Wet hands and forearms, wash with Majola® or Wasa® 2000, rinse well and dry.
Regeneration and Protection of the skin after disinfection:	Luphenil®	Apply on cleandes skin/hands and rub in thoroughly.

Certified and mentioned in the relevant lists.

Date of revision: March 2015

Standard packaging / transport units:

- 20 x 125 ml bottles
- 15 x 250 ml bottles with sprayer (without aerosols)
- 12 x 500 ml bottles
- 12 x 1000 ml bottles
- 2 x 5 litre canisters

Accessories:

Lydos F-wall dispenser for 500 ml bottle

Lydos F-wall dispenser for 1000 ml bottle

**Lysoform Dr. Hans Rosemann GmbH
Kaiser-Wilhelm-Str. 133, D – 12247 Berlin
Tel.: +49-30-77992-0 Telefax: +49-30-77992-219**

0916-10